

Please amend the application as follows:

In the Specification:

Please amend the paragraph beginning on page 1, line 4 as follows:

B1 This application is a ~~continuing application of U.S.S.N.s~~ claims the benefit of the filing date of U.S.S.N. 60/151,987, filed September 1, 1999, and is a continuation in part of U.S.S.N. 09/261,890, filed March 3, 1999, now U.S. Patent No. 6,447,765, which claims the benefit of the filing date of U.S.S.N. 60/076,677, filed March 3, 1998.

Please amend the paragraph beginning on page 8, line 33 as follows:

B2 Figures 7A and B depicts the effect of CD4 cells primed with TGF-beta on allo-cytotoxic T lymphocyte (CTL) activity. The addition of CD4 CD45RA cells that had been cultured for 5 days without stimulators had no effect on CTL activity (result not shown). Culturing these T cells with stimulator cells resulted in modest to moderate suppressive activity. In all experiments, culture of these T cells with TGF-beta 1 ng/ml markedly suppressed, or abolished allo-CTL activity.

Please amend the paragraph beginning on page 9, line 4 as follows:

B3 Figures 8A and 8B demonstrates that regulatory T cells require cell contact to inhibit CTL activity. CD4reg and control CD4+ cells were generated by activating naive CD4+ cells with allogeneic stimulator cells \pm TGF- β as described above. In secondary cultures, CD4reg or control CD4+ cells were either directly mixed with the responder cells or separated from them by a semipermeable membrane using TranswellTM chambers. The Transwells contained CD4+ cells and stimulator cells. The results shown are one of 3 independent experiments.

Please amend the paragraph beginning at page 10, line 27 as follows:

B4 Figures 15A and 15B depicts that repeated stimulation of T cells with a low dose of staphylococcal enterotoxin B (SEB) induces T cells to produce immunosuppressive levels of TGF- β . CD4+ T cells were stimulated with SEB (0.01ng/ml) and irradiated B cells as superantigen presenting cells with or without TGF- β at the times indicated by the arrows. Active TGF- β was measured 2 or 5 days later.

Please amend the paragraph beginning at page 11, line 1 as follows:

B5 Figures 17A-17D shows the effects of SEB on naive (CD45RA+ CD45RO-) CD4+ and CD8+ T cells. The cells were stimulated with SEB every 5th day for a total of three stimulations. The percentages of each T cell subset and the cells expressing the CD25 IL-2 receptor activation marker were determined after each stimulation. Panels Figures 17A and 17C show that if TGF- β 1ng/ml was included in the initial stimulation, CD4+ T cells became the predominant subset in the cultures after repeated stimulation. Panels Figures 17B and 17D show that CD25 expression

35
cont

by SEB stimulated cells decreases by the third stimulation in control cultures. However, CD25 expression remains high if the T cells have been primed with TGF- β .

Please amend the paragraph beginning on page 21, line 15 as follows:

86

Treatment of donor CD8+ Cells ex vivo to suppress an immune attack against blood cells of an unrelated recipient.